



2008-2009 South Carolina Bobwhite Quail Hunter Survey



The South Carolina Department of Natural Resources Small Game Project conducts an annual survey of the state's bobwhite quail (*Colinus virginianus*) hunters. This survey was initiated during the 1987-88 hunting season and was conducted for the 22nd consecutive year during the 2008-09 season. The purpose of the Quail Hunter Survey is to collect quantitative information on hunting success, which aids biologists in tracking quail population trends.

METHODS

A mailing list was generated from individuals who had participated in the Quail Hunter Survey in previous years, applicants from the DNR Public Drawing quail hunts and individuals who responded to news releases or otherwise heard about the survey.

Immediately prior to the opening of the 2008-09 quail hunting season, a total of 205 potential cooperators were provided with materials necessary to participate in the survey. The materials included a pocket-size hunting diary, a data sheet, a postage-paid return envelope, and instructions for the survey. Potential cooperators were also provided with a larger return envelope with tags and asked to collect 10 wings from harvested quail. The wings were used to estimate sex and age ratios for the state's quail population. Funds for printing of the hunting diary booklets were generously provided by the South Carolina State Committee of Quail Unlimited.

Cooperators were instructed to keep accurate records while quail hunting by recording the following information for each hunt: date, county of hunt, number of quail coveys flushed, number of quail bagged, an estimate of average covey size, number of woodcock (*Scolopax minor*) flushed, number of woodcock bagged, time expended on each hunt, and number of rabbits (*Sylvilagus* spp.) seen while quail hunting. Information and comments concerning the overall quality of the season, dog performance, quail food habits, weather, habitat conditions and other observations were also requested. Cooperators were asked to collect 1 wing from each quail harvested (up to 10 per cooperator) and affix a tag with the date, county and sex of the bird to each wing. Survey participants were urged to record information for wild, native bobwhite quail hunts only.

Near the close of the quail season, a reminder postcard was mailed to each potential cooperator encouraging prompt return of the survey forms and wings. Data received were entered on a computer file, proofed and summarized. Data were then analyzed statistically to determine differences by year, county, region, month, or a combination of these variables.

RESULTS AND DISCUSSION

Statewide

Responses were received from 51 individuals, with 44 hunters providing usable data. The total response rate was 21%. Information was obtained from 437 hunts, totaling 1407.5 quail hunting hours. The highest frequency of survey quail hunting activity occurred during the month of January, closely followed by February (Table 1).

Table 1. Quail hunting activity by month, 2008-09 South Carolina Quail Hunter Survey

Month	Number of Hunts	Percentage of Total
November	26	5.9
December	114	26.1
January	152	34.8
February	142	32.5
March	3	0.7
TOTAL	437	100.0

Survey cooperators flushed an average of 0.58 coveys per hour during the 2008-09 season, which was identical to the flush rate in the previous season (Table 2). Quail bagged per hour increased and rabbits jumped per hour increased slightly from last year. The reported average covey size was significantly lower than the previous year. The trend in the coveys per hour index over the 22 years of the survey is illustrated in Figure 1. Quail hunters responding to the survey reported flushing 236 woodcock while quail hunting, for a flush rate of 0.17 woodcock per hour, significantly higher than the 0.02 woodcock flushed per hour in 2007-08. The number of woodcock harvested was 49 during the 2008-09 season, up from 4 woodcock harvested during the 2007-08 season.




Figure 1. Coveys per hour index and trend, South Carolina Quail Hunter Survey, 1988-2009.

Table 2. Statewide South Carolina Quail Hunter Survey results, 1988-2009.

Year of Survey	Coveys Found Per Hour	Quail Bagged Per Hour	Average Covey Size	Rabbits Jumped Per Hour
1987-88	0.76	0.99	12.5	0.21
1988-89	0.72	0.94	12.2	0.19
1989-90	0.80*	1.04	12.9	0.16
1990-91	0.87*	1.07	12.5	0.19*
1991-92	0.65*	0.75*	12.2	0.15*
1992-93	0.66	0.73	12.2	0.15
1993-94	0.67	0.73	11.9	0.22*
1994-95	0.62	0.72	11.8	0.17*
1995-96	0.64	0.76	11.7	0.16
1996-97	0.72*	0.79	12.3	0.18
1997-98	0.63*	0.66*	11.9	0.14
1998-99	0.59	0.62	11.3*	0.16
1999-00	0.42*	0.38*	10.7*	0.14
2000-01	0.58*	0.55*	11.4*	0.13
2001-02	0.62	0.56	11.8	0.17*
2002-03	0.59	0.56	11.2	0.16
2003-04	0.53*	0.49	11.6*	0.13
2004-05	0.54	0.55	11.1*	0.14
2005-06	0.49	0.42*	10.5	0.13
2006-07	0.59*	0.43	11.2*	0.14
2007-08	0.58	0.33*	12.1*	0.15
2008-09	0.58	0.41	10.9*	0.18

*Significantly different from the previous year (P<0.05)

The Quail Hunter Survey also provides information on hunting success by month, region, and county. Hunting success rates as measured by coveys per hour were higher in November and lower in February when compared with the previous season (Fig. 2). Due to a lack of reported hunts in March, 2009, the coveys per hour index was not calculated for that month.




Figure 2. Coveys per hour index by month for the 2007-08 and 2008-09 seasons, South Carolina Quail Hunter Survey.

Regional Comparisons

South Carolina can be divided into 4 broad geographical regions based on habitat type similarities. The Piedmont, which includes the mountain and foothills areas, encompasses about 1/3 of the state. The Midlands extends in a broad band of mostly sandhills habitat on a northeast - southwest axis across the middle of South Carolina. The Northern Coastal Plain or "Pee Dee" and the Southern Coastal Plain or "Low Country" are the other 2 regions.

Hunting effort expended in the 2008-09 Quail Hunter Survey was greatest in the Midlands, accounting for 42% of all hunts (Fig. 3). The coveys per hour index was highest in the Northern Coastal Plain, followed by the Midlands, the Southern Coastal Plain and the Piedmont (Table 3). Quail harvest rate was higher in the Northern Coastal Plain than in the other three regions. The reported average size of quail coveys ranged from 8.8 birds per covey in the Piedmont to 11.2 birds per covey in the Midlands. The rabbit jumping rate by quail hunters was highest in the Northern Coastal Plain and lowest in the Southern Coastal Plain.




Figure 3. Percentage of total hunts occurring in each region, South Carolina Quail Hunter Survey, 2008-09.

Table 3. Results by region, South Carolina Quail Hunter Survey, 2008-09.

Region	Coveys Found Per Hour*	Quail Bagged Per Hour*	Average Covey Size*	Rabbits Jumped Per Hour*
N. Coastal Plain	0.93A	0.60A	10.7A	0.24A
Midlands	0.55B	0.43AB	11.2A	0.18AB
S. Coastal Plain	0.35C	0.26BC	10.8A	0.04B
Piedmont	0.12D	0.07C	8.8B	0.10AB

*Numbers followed by the same letter within the same column are not significantly different (P > 0.05).

County Results

The top 5 counties for each of 4 indices are listed in Table 4. A minimum of 15 sample hunts had to occur in a particular county to be included in this ranking. Clarendon County claimed the top spot in the coveys per hour index at 1.63, followed by Sumter (0.91) and Lee (0.69) counties. Clarendon County also led the state in quail harvested per hour at 1.11, followed by Sumter (0.77) and Darlington (0.60) counties.

The top five counties for quail hunting lie in the Northern Coastal Plain Region or Midlands Region. County results should be interpreted cautiously due to the relatively low number of hunts upon which they are based. Nevertheless, Northern Coastal Plain and Midlands counties have proven to be the better areas for South Carolina quail hunting (Fig. 4).

Table 4. Ranking of the top counties for indices measured in the South Carolina Quail Hunter Survey, 2008-09.

Rank*	Coveys Found Per Hour	Quail Harvested Per Hour	Average Covey Size	Rabbits Jumped Per Hour
1	Clarendon (1.63)	Clarendon (1.11)	Calhoun (12.46)	Horry (0.40)
2	Sumter (0.91)	Sumter (0.77)	Darlington (12.33)	Lee (0.39)
3	Lee (0.69)	Darlington (0.60)	Lee (11.40)	Florence (0.15)
4	Horry (0.67)	Lee (0.45)	Sumter (11.33)	Laurens (0.14)
5	Darlington (0.53)	Horry (0.35)	Chesterfield (10.65)	Darlington (0.10)

*Counties with a sample size of less than 15 hunts not included.




Figure 4. Coveys per hour index by region for the 2007-08 and 2008-09 seasons, South Carolina Quail Hunter Survey.

Quail Wing Data

All potential Quail Hunter Survey cooperators were asked to provide wings from harvested birds. Twenty-six survey participants provided usable wings, for a response rate of 59% of those providing hunting data. Age information was obtained from 164 wings collected during the 2008-09 season, and cooperators provided the sex of the birds from which 147 wings were taken.

A higher percentage of males is believed to be normal for fall quail populations because of the hen's greater susceptibility to predation and physical stress due to egg laying and incubation during the nesting season. The sex ratio obtained from quail wing samples during the 2008-09 season was 56% males to 44% females (Fig. 5). Overall, the percentage of juveniles in the harvest was 84%, versus 16% adults (Fig. 5). The age ratio of 5.31 juveniles per adult was significantly higher than the age ratio of 2.46 juveniles per adult recorded in the 2007-08 survey. This increased age ratio is likely a reflection of better nesting and brood rearing conditions during the 2008 nesting season following drought conditions in 2007. Age ratios in quail of 5-6 juveniles per adult are considered good and 7-8 juveniles per adult excellent.




Figure 5. Quail sex and age distribution from sample wings, South Carolina Quail Hunter Survey, 2008-09.

Comments From Cooperators

Only 7% of those responding rated the 2008-09 season as good, while 37% gave the season a fair rating, and 56% a poor rating (Fig. 6). Dog performance was rated good by 56% and fair by 37% of cooperators (Fig. 6).




Figure 6. Ratings of season quality and dog performance by cooperators in the South Carolina Quail Hunter Survey, 2008-09.

Cooperator comments provide insight into general habitat and hunting conditions and predominant food items utilized by quail during fall and winter. Commonly submitted comments on the 2008-09 quail season are listed below. Number of similar comments received is noted in parentheses.

-  Season was poor/worse than last year (6).
-  More birds than past few years (3).
-  Habitat appears good (4).

Typically, a basic food habits analysis is provided as part of this report. However, for the 2008-09 season, cooperators did not provide enough information regarding quail food habits to allow a summary.

Plans are underway to continue the Quail Hunter Survey for the 2009-10 season. Efforts are ongoing to increase the number of cooperators and all avid quail hunters are encouraged to participate. Contact the SCDNR Small Game Project, P.O. Box 167, Columbia, SC 29202 (telephone 803-734-3609) to be included in the survey.