

2024 Bobwhite Quail Whistling Cock Census

The Bobwhite Quail Whistling Cock Census was conducted for the 46th consecutive year in 2024. Seventy-six routes were sampled in 2024, resulting in sixty-eight routes which yielded data comparable to the previous year. Approximate locations of survey routes are shown in Figure 1.




Figure 1. South Carolina Bobwhite Quail Whistling Cock Census route locations - 1979 - 2024.

Methods

Census routes are sampled between 15 June and 10 July, believed to be the period of maximum bobwhite whistling in the Southeast (Rosene 1969). Surveys began at official sunrise, with routes consisting of 12 stops at ½ mile intervals. The observer listened for 8 minutes at each stop and recorded the number of calling males heard. Weather conditions (i.e., cloud cover, wind speed, # add in any others) were recorded at each stop. Operation of routes was discouraged on rainy or windy days. Number of calling males at each stop were summed to get a total for the route. Based on an extensive analysis of historical Whistling Cock Census data, the survey protocol was changed in 2003 from a two-day survey to a one-day survey. The afternoon listening period was also eliminated as an option in the survey protocol.

As in previous years, data were analyzed using a 2-tailed paired *t*-test for equal sample sizes (Steel and Torrie 1980). Between-year comparisons were conducted using all routes that were run in both 2024 and 2023. The 2024 mean was also compared with the long-term average using a 2-tailed paired *t*-test.

Results and Discussion

Approximately fifty-eight observers surveyed seventy-six routes in 2024. Sixty-eight of these routes were used in year-to-year comparisons. A total of 326 whistling cocks were recorded on the sixty-eight routes for an average of 4.8 (SE = 0.94) birds/route. For all routes sampled in 2024, a total of 366 whistling cocks were recorded with an average number of calling birds of 4.8 (SE = 0.86) birds/route (Table 1).

The 2024 mean call count index did not significantly differ ($P > 0.05$) from the 2023 index. The 2024 index was significantly less than ($P < 0.05$) the long-term average for the 46 years of the census (11.2 birds/route, $N=2373$, Fig. 2).

Table 1. Summary of South Carolina Bobwhite Quail Whistling Cock Survey Census, 1979-2024.

Year	Number of Routes	Calling Males/Route (Average)
1979	18	33.6
1980	19	30.5
1981	26	33.8
1982	28	32.0
1983	27	40.1
1984	28	29.3
1985	28	31.6
1986	30	29.1
1987	28	33.4
1988	28	31.8
1989	38	22.0
1990	39	21.5
1991	40	16.8
1992	43	17.9
1993	44	17.9
1994	47	15.5
1995	51	15.0
1996	53	16.7
1997	52	13.3
1998	53	13.5
1999	56	10.0
2000	57	9.3
2001	56	11.1
2002	59	12.6
2003	60	10.3
2004	61	11.6
2005	61	12.7
2006	62	11.7
2007	63	8.0
2008	63	8.6
2009	66	9.9
2010	67	9.3
2011	68	8.4
2012	64	8.2
2013	62	8.0
2014	47	5.1
2015	59	3.6
2016	63	4.5
2017	62	5.2
2018	69	6.3
2019	72	5.4
2020	66	6.2
2021	74	6.7
2022	74	5.9
2023	68	5.9
2024	76	4.8
46-year average	52	11.2




Figure 2: Average Bobwhite Quail Whistling Cock Census call count (birds/route) in South Carolina 1979-2024, trendline, and long-term average for the period.

Quail call counts in 2024 were higher on twenty-three routes (34 %) and lower on thirty-four routes (50 %) compared to 2023. Statewide, there was 75 less birds recorded in 2024 than were recorded on the same routes in 2023, a decrease of 19%.

Because of natural mortality, breeding populations of quail are never high relative to early fall populations. Extreme weather conditions during the nesting season and land-use changes can further depress quail numbers.

Whistling Cock Survey data are used in conjunction with Quail Brood Survey data, Quail Hunter Survey data, and Fall Covey Count data to assess the population status of quail statewide as well as the effects of land use change and other factors, such as weather, on the statewide quail population.

Land-use changes surrounding routes is the most difficult variable to qualify. When routes for the Whistling Cock Census were established in 1979, cooperators were instructed to locate census lines in areas where good quail populations existed. Routes are not moved to new locations unless access is limited or noise disturbance (e.g., road traffic) makes counting calling birds impossible. Many routes which were established through mixed forest and farmlands are now surrounded by pine plantations. Therefore, a dramatic decrease in the suitability of the habitat surrounding survey routes is thought to have severely depressed the numbers of calling birds on many routes. While large-scale (e.g. state level) habitat changes are obvious and are easily documented, it is much more difficult to quantify habitat changes at the call count route or

point level. Analysis of fine-scale habitat components along survey routes should be conducted to validate the effects of habitat change on local quail populations.

It is important to note that the full complement of routes were established to document year-to-year and long-term population changes. Thus, populations are expected to decrease for a period of time and should rebound if the habitat reverts to a more suitable type.

The Bobwhite Quail Whistling Cock Census should be continued in order to monitor changes in habitat conditions and the resultant quail population response.

Literature Cited

- Rosene, W., Jr. 1969. The Bobwhite Quail: Its life and management. Rutgers University Press, New Brunswick, N. J. 418 pp.
- Steel, R. G. D. and J. H. Torrie. 1980. Principles and procedures of statistics: A biometrical approach. McGraw Hill Book Company, New York. 633 pp.